

CEFQUINOX® LC

Intramammary Suspension

Fourth Generation Cephalosporin Against All
Relevant Bacteria Causing Bovine Mastitis

Agrovet MARKET



Formulation

Each 8 g syringe contains: Cefquinome (as sulfate) 75 mg, excipients q.s.ad. 8 g.

Indications

For the treatment of clinical and subclinical mastitis in lactating cows caused by bacteria sensitive to cefquinome, namely *Staphylococcus aureus*, *Streptococcus uberis*, *Streptococcus dysgalactiae*, *Escherichia coli* (and other enterobacteria).

Dosage and Administration

Cefquinox® LC is administered intramammary. One syringe per affected quarter every 12 hours for 3 successive milkings. Milk the affected quarters and discard the milk. After thoroughly cleaning and disinfecting the teat and teat sphincter, gently infuse the contents of one syringe into the teat canal of each affected quarter. Disperse the product by gently massaging the teat and udder of the affected animal. The contents of one syringe should be gently infused into the teat of the infected quarter every 12 hours after each of three successive milkings. Take precautions to avoid contamination of the syringe cannula.

Commercial Presentation

Container x 48 syringes x 3 g,

SENASA Registry

Perú: F.082.005.I.00996.



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.