

FLORAVIVA® 12 CEPAS SOFT CHEWS MINI & SMALL

Tablet Oral

Protective symbiotic, improver and recuperator of intestinal balance Live Active Cultures - Gut Health



Formulation

Each 2 g chewable tablet contains: Probiotics (12 cultures) 10 x 10⁸ CFU, Active Organic Dry Yeast Extract 100 mg, -glucans 40 mg, Fructooligosaccharides (FOS) 40 mg, Mannan oligosaccharides (MOS) 40 mg, Lactoferrin 20 mg, Yucca schidigera extract 20 mg, Glutamic Acid 16 mg, L-Threonine 12 mg, Vitamin A (Retinol Palmitate) 5,000 IU, Vitamin D3 (as cholecalciferol) 1,000 IU, Vitamin E (as alpha tocopheryl acetate) 8 mg, Vitamin K3 (as menadione sodium bisulfite) 500 ug, Vitamin B1 (Thiamine hydrochloride) 8 mg, Vitamin B2 (Riboflavin 5 Sodium Phosphate) 8 mg, Vitamin B3 (Nicotinamide) 20 mg, Vitamin B5 (Calcium Pantothenate) 30 mg, Vitamin B6 HCl (Pyridoxine) 4 mg, Vitamin B7 (Biotin) 40 ug, Vitamin B9 (Folic Acid) 500 ug, Vitamin B12 (Cyanocobalamin) 20 ug, Vitamin C (as sodium ascorbate) 50 mg, Copper (copper chelate-AHM2) 1,020 ug, Zinc (zinc chelate-AHM2) 1,292 ug, Manganese (manganese chelate-AHM2) 1,050 ug, L-Selenomethionine 24 ug, Methionine (AHM3) 108 mg, Excipients and flavorings q.s.ad. 2 g.

Indications

Auxiliary in the recomposition and stabilization of the intestinal flora and physiological digestion, as well as improving immunity and protecting the intestinal structure and functionality in cases of:- Diarrhea or digestive disorders, poor fecal quality (soft stools).- Acute enteritis- Allergies or intolerance to food.- Poor fecal quality, flatulence,- Antibiotic, chemotherapy or antiparasitic therapy.- Birth and/or weaning.- Stress conditions: postoperative, childbirth, excessive heat or cold, travel, vaccination, heat, exhibitions, mating.- Infectious diseases (parasitic, viral or bacterial). Strengthens the immune system.- Changes in diet or alterations in the type of food.- Poisoning and/or poisoning.- Neonates and young growing animals.- Old age animals.- Intestinal dysbiosis (imbalance of the intestinal flora).- Pancreatic problems, malabsorption syndrome and chronic liver disorders.

Dosage and Administration

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso. It is administered orally, according to the size of the pet. Administer 1 to 2 times a day according to the following dosage: - Puppies (up to 3 kg) and kittens: ½ tablet - Miniature