

ULTRAMETRIN® 600

Emulsifiable Concentrate

Pyrethroid-phosphorous potentialized association







Formulation

Cypermethrin 100 mg, Ethion 400 mg, Piperonyl butoxide 100 mg, excipients q.s. ad 1 mL.

Indications

The product combats the common cattle tick Boophilus microplus and other tick species. It also controls a number of biting insects like horn flies, stable flies, mosquitos and horse flies, as well as mites, lice and fleas, among others. Its residual action lasts up to 4 weeks.

Dosage and Administration

For immersion baths: Foot bath: 1 L/1000 L of fresh or salty water. Replenishing and restrengthening the solution: 1.5 L/1000 L of water. Both operations to be performed when the level in the bath is reduced in 10% of its original level. Dry restrengthening: 0.5 L/1000 L of water consumed in the bath. This is done in order to not having to replenish the water consumed. For sprinkling animals: 1 L/1000 L of water (apply the final sprinkling emulsion using approximately 1 L/100 kg of b.w.). For spraying in buildings and facilities: 1 mL/1 to 2 L of water (1 L of the final emulsion per 10 to 15 m2 of surface to be treated).

Commercial Presentation

Bottle x 20 mL, Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Bottle x 1 L,

SENASA Registry

SENASA Perú: F.07.08.N.0327



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.